

On Combinatorial Problem Concerning Partitions of a Box into Boxes

Apoloniusz Tyszk


Abstract

We consider partitions of n -dimensional boxes in $\mathbb{R}^n$, $n \geq 2$, into a finite number of boxes with pairwise disjoint interiors. We study sets $X \subseteq (0, \infty)$ with the Property (W_n) : for every n -dimensional box P and every partition of P , if each constituent box has one side with the length belonging to X , then the length of one side of P belongs to X . We prove that the set $X \subseteq (0, \infty)$ has Property (W_n) if and only if X is closed with respect to the operations: $x+y$ and $x+y+z-2\min(x, y, z)$.

We consider partitions of n -dimensional boxes in $\mathbb{R}^n$, $n \geq 2$, into a finite number of n -dimensional boxes with pairwise disjoint interiors. We study sets $X \subseteq (0, \infty)$ with the Property (W_n) : for every n -dimensional box P and every partition of P , if each constituent box has one side with the length belonging to X , then the length of one side of P belongs to X .

Theorem (see: [5] for groups, [4] for generalizations of groups, [1]–[3] for earlier results). *The set $X \subseteq (0, \infty)$ has Property (W_n) if and only if X is closed with respect to the operations: $x + y$ and $x + y + z - 2\min(x, y, z)$.*

Proof. Necessity: We can assume that $x > z$, $y > z$. Case $n = 2$: the result follows from the following partitions of a square:



Case $n > 2$: we multiply the first square by $[0, x + y]^{n-1}$, the second square by $[0, x + y - z]^{n-1}$. Sufficiency: we first prove (see Eulerian path method in [5] by Michael S. Paterson, Univ. of Warwick, Coventry, England) that if each constituent box from the partition of the box P has one side with the length belonging to X , then there exist points $Y_1, Y_2, \dots, Y_m$ lying on the one side of P such that this side is equal to the segment $\overline{Y_1 Y_m}$ and every distance $|Y_i Y_{i+1}|$ belongs to X . We choose the cartesian coordinate system with axes $x_j (j = 1, 2, \dots, n)$ which are parallel to the sides of the box P . For each constituent box P_k we choose $c(k) \in \{1, 2, \dots, n\}$ such that the length of P_k in direction $x_{c(k)}$ belongs to X . We define an undirected graph G in the following way: as a vertex set we put the set of all vertices of constituent boxes, as an edge set we put the set of all pairs (s, P_k) , where s is a side of P_k lying in the direction $x_{c(k)}$. Each vertex of G (except the vertices of the box P which lie on 1 edge) is a vertex of an even number of constituent boxes, hence it lies on an even number of edges of G . Thus a walk away along edges that begins at one vertex of P and does not repeat any edges will not terminate until it hits another vertex of P . This observation produce points $Y_1, Y_2, \dots, Y_m$.

Now we are ready to prove that $|Y_1 Y_m| \in X$. It suffices to prove that if there exist m points with the required property (for $m > 3$), then there exist $m - 1$ or $m - 2$ points with this property. We can assume $Y_1, Y_2, \dots, Y_m$ are different and for every i such that $1 < i < m$, $Y_i \notin \overline{Y_{i-1} Y_{i+1}}$, hence $Y_3 \in \overline{Y_1 Y_2}$ and $Y_{m-2} \in \overline{Y_{m-1} Y_m}$, so $|Y_2 Y_3| < |Y_1 Y_2|$ and $|Y_{m-1} Y_m| > |Y_{m-2} Y_{m-1}|$. We choose the smallest $i > 2$ such that $|Y_i Y_{i+1}| > |Y_{i-1} Y_i|$. From this choice of i we have $|Y_{i-1} Y_i| < |Y_{i-2} Y_{i-1}|$, so

$$\bullet \xrightarrow{Y_{i-2}} \bullet \xrightarrow{Y_i} \bullet \xrightarrow{Y_{i-1}} \bullet \xrightarrow{Y_{i+1}}$$

$$\begin{aligned} |Y_{i-2} Y_{i+1}| &= |Y_{i-2} Y_{i-1}| - |Y_{i-1} Y_i| + |Y_i Y_{i+1}| = \\ &= |Y_{i-2} Y_{i-1}| + |Y_{i-1} Y_i| + |Y_i Y_{i+1}| - 2 \min(|Y_{i-2} Y_{i-1}|, |Y_{i-1} Y_i|, |Y_i Y_{i+1}|). \end{aligned}$$

X is closed with respect to the operation $x + y + z - 2 \min(x, y, z)$, hence $|Y_{i-2} Y_{i+1}| \in X$ and we can reduce the number of points to $m - 2$, this ends the proof of sufficiency.

Our proof is now complete. □

References

- [1] N. G. de Bruijn, *Filling boxes with bricks*, Am. Math. Monthly **76** (1969), 37–40.
- [2] M. Dehn, *Über Zerlegung von Rechtecken in Rechtecke*, Math. Ann. **57** (1903), 314–332.
- [3] R. Johnsonbaugh, *De Bruijn's packing problem*, in *Two-year college mathematics readings*, (ed. W. Page), Math. Ass. Am., 1981, pp. 258–262.
- [4] A. Tyszką, *On some combinatorial problems concerning partitions of a box*, J. Natur. Geom. **5** (1994), 1–9,
<http://www.cyf-kr.edu.pl/~rttyszka/jnatgeom1994.doc>
- [5] S. Wagon, *Fourteen proofs of a result about tiling a rectangle*, Am. Math. Monthly **94** (1987), 601–617.

Apoloniusz Tyszką
Technical Faculty
Hugo Kołłątaj University
Balicka 116B, 30-149 Kraków, Poland
E-mail address: rttyszka@cyf-kr.edu.pl